arXiv:1412.8358v3 [math.CO] 31 Aug 2015

Odd graph and its applications on the strong edge coloring

Tao Wang* Xiaodan Zhao
Institute of Applied Mathematics
Henan University, Kaifeng, 475004, P. R. China

August 15, 2015

Abstract

A strong edge coloring of a graph is a proper edge coloring in which every color class is an induced matching. The strong chromatic index $\chi_s'(G)$ of a graph G is the minimum number of colors in a strong edge coloring of G. Let $\Delta \geqslant 4$ be an integer. In this note, we study the properties of the odd graphs, and show that every planar graph with maximum degree at most Δ and girth at least $10\Delta-4$ has a strong edge coloring with $2\Delta-1$ colors. In addition, we prove that if G is a graph with girth at least $2\Delta-1$ and $\mathrm{mad}(G) < 2+\frac{1}{3\Delta-2}$, where $\Delta \geqslant 4$, then $\chi_s'(G) \leqslant 2\Delta-1$; if G is a subcubic graph with girth at least 8 and $\mathrm{mad}(G) < 2+\frac{2}{23}$, then $\chi_s'(G) \leqslant 5$.

1 Introduction

All graphs considered in this paper are simple, finite and undirected. A *strong edge coloring* of a graph is a proper edge coloring in which every color class is an induced matching. The *strong chromatic index* $\chi_s'(G)$ of a graph G is the minimum number of colors in a strong edge coloring of G. Fouquet and Jolivet [13, 14] introduced the notion of strong edge coloring for the radio networks and their frequencies assignment problem. More details on the applications of strong edge coloring can be found in [2, 23, 25].

Greedy algorithm provides an upper bound $2\Delta(\Delta-1)+1$ on the strong chromatic index, where Δ is the maximum degree of the graph. In 1985, Erdős and Nešetřil constructed graphs with strong chromatic index $\frac{5}{4}\Delta^2$ when Δ is even, $\frac{1}{4}(5\Delta^2-2\Delta+1)$ when Δ is odd. Inspired by their construction, they proposed the following strong edge coloring conjecture during a seminar in Prague.

Conjecture 1 ([10])**.** If G is a graph with maximum degree Δ, then

$$\chi_s'(G) \leqslant \begin{cases} \frac{5}{4}\Delta^2, & \text{if } \Delta \text{ is even,} \\ \frac{1}{4}(5\Delta^2-2\Delta+1), & \text{if } \Delta \text{ is odd.} \end{cases} \quad \square$$

In 1997, Molloy and Reed [20] gave the first general bound by using probabilistic techniques, showing that if Δ is large enough, then every graph with maximum degree Δ has a strong edge coloring with $1.998\Delta^2$ colors. Recently, Bruhn and Joos [4] announced an improvement from $1.998\Delta^2$ to $1.93\Delta^2$.

In 1990, Faudree, Schelp, Gyárfás and Z. Tuza [12] proposed various conjectures on subcubic graphs (graphs with maximum degree at most three).

Conjecture 2. Let G be a subcubic graph.

1. The strong chromatic index is at most 10.
2. If G is bipartite, then $\chi_s'(G) \leqslant 9$.
3. If G is planar, then $\chi_s'(G) \leqslant 9$.
4. If G is bipartite and the degree sum of every edge is at most 5, then $\chi_s'(G) \leqslant 6$.

*Corresponding author: wangtao@henu.edu.cn

5. If G is bipartite with girth at least 6, then $\chi'_s(G) \leqslant 7$.

6. If G is bipartite and its girth is large, then $\chi'_s(G) \leqslant 5$.

Conjecture 2.1, the case when $\Delta \leqslant 3$ in Conjecture 1, was confirmed independently by Andersen [1] and Horák et al. [16]. Faudree, Gyárfás, Schelp and Z. Tuza [11] conjectured that $\chi'_s(G) \leqslant \Delta(G)^2$ if G is a bipartite graph. In 1993, Steger and Yu [28] confirmed Conjecture 2.2, which is the first nontrivial case of Faudree's conjecture on bipartite graph. Very recently, Kostochka et al. [17] proved Conjecture 2.3; in fact, they proved the result for subcubic planar multigraph without loops (multiple edges are allowed). Recently, Conjecture 2.4 was verified by Wu and Lin [30], and it was generalized by Nakprasit [21, 22]. The strong chromatic index of some classes of graphs have been extensively studied, such as degenerate graphs [7, 9, 29, 31], Halin graphs [5, 18, 19, 26, 27].

Regarding the planar graphs, Faudree et al. [12] established the following upper bound using the Four Color Theorem and Vizing's Theorem.

Theorem 1.1. Every planar graph with maximum degree Δ has a strong edge coloring with $4\Delta + 4$ colors.

Borodin and Ivanova [3] showed that if G is a planar graph with girth at least $40\lfloor\frac{\Delta}{2}\rfloor + 1$, then $\chi'_s(G) \leqslant 2\Delta - 1$. Chang, Montassier, Pêcher and A. Raspaud [6] improved the lower bound on the girth as following.

Theorem 1.2. Let $\Delta \geqslant 4$ be an integer. If G is a planar graph with maximum degree at most Δ and girth at least $10\Delta + 46$, then $\chi'_s(G) \leqslant 2\Delta - 1$.

Recently, Guo et al. [15] improve the girth to $10\Delta + 26$. In this note, we give a better result on the odd graph by following Chang's method, and then improve the lower bound on the girth to $10\Delta - 4$ when $\Delta \geqslant 4$. In addition, we give two results with the condition on the maximum average degree. The maximum average degree $\mathrm{mad}(G)$ of a graph G is

$$\mathrm{mad}(G) = \max_{H \subseteq G} \left\{ \frac{2|E(H)|}{|V(H)|} \right\}.$$

2 Odd graph

The odd graph O_n has one vertex for each of the $(n-1)$-element subsets of a $(2n-1)$-element set. Two vertices are adjacent if and only if the corresponding subsets are disjoint. We use v to denote a vertex in the odd graph, and use $[v]$ to denote the corresponding subset. By the definition, every edge uv can be labeled with a unique element which is not contained in $[u]$ or $[v]$. By the particular structure of odd graph, the labels on the edges induce a strong edge coloring of the odd graph. We will use this property in the proof of Theorem 3.3. A *walk* is a sequence of edges such that every two consecutive edges are adjacent. A *special walk* is a walk whose every two consecutive edges are distinct. A *k-special walk* is a special walk of length k.

2.1 Structural results

Theorem 2.1. If the distance between u and v is two, then $[u]$ and $[v]$ have exactly one different element. Thus, for two vertices u and v, if $[u]$ and $[v]$ have exactly k different elements, then they can be reached from each other in $2k$ steps and this path is a shortest even path.

Theorem 2.2. If $w_1w_2w_3$ is a 2-path with w_1w_2 and w_2w_3 are labeled with x and y respectively, then $[w_3]$ is obtained from $[w_1]$ by replacing y with x.

In the following, we use $\underbrace{x, y}$ to denote the labels x and y on two consecutive edges.

Theorem 2.3. Every 3-path $v_1v_2v_3v_4$ is contained in a 6-cycle.

Proof. Note that the sets corresponding to v_1 and v_3 have only one different element. So we may assume that $[v_1] = X \cup \{x_1\}$ and $[v_3] = X \cup \{x_2\}$, where X is an $(n-2)$-subsets of $\{1, 2, \ldots, 2n-1\}$. Thus, we can obtain $[v_2] = \{1, 2, \ldots, 2n-1\} \backslash (X \cup \{x_1, x_2\})$. Without loss of generality, we assume that $[v_4] = \{1, 2, \ldots, 2n-1\} \backslash (X \cup \{x_2, x_3\})$. It is easy to check that the path $v_1v_2v_3v_4$ is contained in a 6-cycle $v_1v_2v_3v_4v_5v_6$, where v_5 and v_6 are labeled with $X \cup \{x_3\}$ and $\{1, 2, \ldots, 2n-1\} \backslash (X \cup \{x_1, x_3\})$. □

Theorem 2.4. Given two vertices w_0, w_{2n} (not necessarily distinct) in O_n ($n \geqslant 4$) with $\lambda_1 \notin [w_0]$ and $\lambda_2 \notin [w_{2n}]$, there exists a $2n$-special walk from w_0 to w_{2n} such that its first edge is labeled with λ_1 and the last edge is labeled with λ_2.

Proof. We may assume that $[w_0] = \{x_1, x_2, \ldots, x_k\} \cup \{y_{k+1}, y_{k+2}, \ldots, y_{n-1}\}$, $[w_{2n}] = \{x_1, x_2, \ldots, x_k\} \cup \{z_{k+1}, z_{k+2}, \ldots, z_{n-1}\}$ and $[w_0] \cup [w_{2n}] \cup \{s_1, s_2, \ldots, s_{k+1}\} = \{1, 2, \ldots, 2n-1\}$. By symmetry, we divide the proof into three cases.

Case 1. $\lambda_1 \in \{z_{k+1}, z_{k+2}, \ldots, z_{n-1}\}$ and $\lambda_2 \in \{y_{k+1}, y_{k+2}, \ldots, y_{n-1}\}$.

It is obvious that $k \neq n-1$. By symmetry, we may assume that $\lambda_1 = z_{k+1}$ and $\lambda_2 = y_{n-1}$.

Subcase 1.1. $k \leqslant n-4$.

In the odd graph O_n, we can find a special walk from w_0 to w_{2n}, whose edges are labeled with $\underline{z_{k+1}, y_{k+1}}$, $\underline{z_{k+2}, y_{k+2}}$, $\ldots$, $\underline{z_{n-3}, y_{n-3}}$, $\underline{s_1, y_{n-2}}$, $\underline{s_2, s_1}$, $\ldots$, $\underline{s_{k+1}, s_k}$, $\underline{z_{n-2}, s_{k+1}}$, $\underline{z_{n-1}, y_{n-1}}$.

Subcase 1.2. $k = n-3$.

In the odd graph O_n, we can find a special walk from w_0 to w_{2n}, whose edges are labeled with $\underline{z_{n-2}, x_1}$, $\underline{s_1, y_{n-2}}$, $\underline{s_2, s_1}$, $\underline{s_3, s_2}$, $\ldots$, $\underline{s_k, s_{k-1}}$, $\underline{x_1, s_k}$, $\underline{z_{n-1}, y_{n-1}}$. (It needs n at least 4, otherwise x_1 does not exist.)

Subcase 1.3. $k = n-2$.

In the odd graph O_n, we can find a special walk from w_0 to w_{2n}, whose edges are labeled with $\underline{z_{n-1}, y_{n-1}}$, $\underline{s_1, x_1}$, $\underline{s_2, s_1}$, $\underline{s_3, s_2}$, $\ldots$, $\underline{s_{n-3}, s_{n-4}}$, $\underline{y_{n-1}, s_{n-3}}$, $\underline{x_1, y_{n-1}}$.

Case 2. $\lambda_1 \in \{z_{k+1}, z_{k+2}, \ldots, z_{n-1}\}$ and $\lambda_2 \in \{s_1, s_2, \ldots, s_{k+1}\}$.

It is obvious that $k \neq n-1$. By symmetry, we may assume that $\lambda_1 = z_{k+1}$ and $\lambda_2 = s_{k+1}$.

If $k = 0$, then we can find a special walk from w_0 to w_{2n}, whose edges are labeled with $\underline{z_1, y_1}$, $\underline{z_2, y_2}$, $\ldots$, $\underline{z_{n-2}, y_{n-2}}$, $\underline{s_1, y_{n-1}}$, $\underline{z_{n-1}, s_1}$.

If $k \in \{1, 2, \ldots, n-2\}$, then we can find a special walk from w_0 to w_{2n}, whose edges are labeled with $\underline{z_{k+1}, y_{k+1}}$, $\underline{z_{k+2}, y_{k+2}}$, $\ldots$, $\underline{z_{n-2}, y_{n-2}}$, $\underline{s_1, y_{n-1}}$, $\underline{s_2, s_1}$, $\underline{s_3, s_2}$, $\ldots$, $\underline{s_{k+1}, s_k}$, $\underline{z_{n-1}, s_{k+1}}$.

Case 3. $\lambda_1 \in \{s_1, s_2, \ldots, s_{k+1}\}$ and $\lambda_2 \in \{s_1, s_2, \ldots, s_{k+1}\}$.

Subcase 3.1. $\lambda_1 = \lambda_2 = s_1$.

If $k = 0$, then we can find a special walk whose edges are labeled with $\underline{s_1, y_1}$, $\underline{z_2, y_2}$, $\underline{z_3, y_3}$, $\ldots$, $\underline{z_{n-1}, y_{n-1}}$, $\underline{z_1, s_1}$.

If $k = 1$, then we can find a special walk whose edges are labeled with $\underline{s_1, x_1}$, $\underline{z_2, y_2}$, $\underline{z_3, y_3}$, $\ldots$, $\underline{z_{n-1}, y_{n-1}}$, $\underline{x_1, s_1}$.

If $k = 2$, then we can find a special walk whose edges are labeled with $\underline{s_1, x_1}$, $\underline{s_3, y_3}$, $\underline{z_3, s_3}$, $\underline{z_4, y_4}$, $\ldots$, $\underline{z_{n-1}, y_{n-1}}$, $\underline{x_1, s_1}$.

If $k \in \{3, \ldots, n-2\}$, then we can find a special walk whose edges are labeled with $\underline{s_1, x_1}$, $\underline{s_3, y_{k+1}}$, $\underline{s_4, s_3}$, $\underline{s_5, s_4}$, $\ldots$, $\underline{s_{k+1}, s_k}$, $\underline{z_{k+1}, s_{k+1}}$, $\underline{z_{k+2}, y_{k+2}}$, $\ldots$, $\underline{z_{n-1}, y_{n-1}}$, $\underline{x_1, s_1}$.

If $k = n-1$, then we can find a special walk whose edges are labeled with $s_1, \underline{x_2, s_2}, \underline{x_3, x_2}, \underline{x_4, x_3}, \ldots, \underline{x_k, x_{k-1}}, \underline{s_2, x_k}, s_1$.

Subcase 3.2. $\lambda_1 = s_1$ and $\lambda_2 = s_{k+1}$. It follows that $k \neq 0$.

If $k = 1$, then we can find a special walk whose edges are labeled with $\underline{s_1, y_2}$, $\underline{s_2, s_1}$, $\underline{z_3, y_3}$, $\underline{z_4, y_4}$, $\underline{z_5, y_5}$, $\ldots$, $\underline{z_{n-1}, y_{n-1}}$, $\underline{z_2, s_2}$.

If $k = 2$, then we can find a special walk whose edges are labeled with $\underline{s_1, y_3}$, $\underline{s_2, s_1}$, $\underline{s_3, s_2}$, $\underline{z_4, y_4}$, $\underline{z_5, y_5}$, $\ldots$, $\underline{z_{n-1}, y_{n-1}}$, $\underline{z_3, s_3}$.

If $k \in \{3, 4, \ldots, n-2\}$, then we can find a special walk whose edges are labeled with $\underline{s_1, y_{k+1}}$, $\underline{s_2, s_1}$, $\underline{s_3, s_2}$, $\underline{s_4, s_3}$, $\underline{s_5, s_4}$, $\ldots$, $\underline{s_{k+1}, s_k}$, $\underline{z_{k+2}, y_{k+2}}$, $\underline{z_{k+3}, y_{k+3}}$, $\ldots$, $\underline{z_{n-1}, y_{n-1}}$, $\underline{z_{k+1}, s_{k+1}}$.

If $k = n-1$, then we can find a special walk whose edges are labeled with $\underbrace{s_2, x_2}$, $\underbrace{s_3, s_2}$, $\underbrace{s_4, s_3}$, $\underbrace{s_5, s_4}$, $\underbrace{s_6, s_5}$, $\ldots$, $\underbrace{s_k, s_{k-1}}$, $\underbrace{s_{k+1}, s_k}$, $\underbrace{x_2, s_{k+1}}$. This completes the proof of the theorem. □

Note that we restrict $n \geqslant 4$ in the above theorem since the result is not true for $n = 3$, see the next subsection. For $n = 3$, we have the following result.

Theorem 2.5. Given two vertices w_0, w_9 (not necessarily distinct) in O_3 with $\lambda_1 \notin [w_0]$ and $\lambda_2 \notin [w_9]$, there exists a 9-special walk from w_0 to w_9 such that its first edge is labeled with λ_1 and the last edge is labeled with λ_2.

Proof. By symmetry, we divide the proof into three cases in terms of the size of intersection of $[w_0]$ and $[w_9]$, and each case is divided into some subcases.

Case 1. Assume that $[w_0] = \{y_1, y_2\}$, $[w_9] = \{z_1, z_2\}$ and $\{y_1, y_2, z_1, z_2, s_1\} = \{1, 2, 3, 4, 5\}$.

When $\lambda_1 = z_1$ and $\lambda_2 = y_1$, we can find a special walk whose edges are labeled with $\underbrace{z_1, y_1}, \underbrace{z_2, y_2}, \underbrace{s_1, z_1}, \underbrace{y_2, z_2}, y_1$.

When $\lambda_1 = z_1$ and $\lambda_2 = s_1$, we can find a special walk whose edges are labeled with $\underbrace{z_1, y_1}, \underbrace{z_2, y_2}, \underbrace{y_1, z_1}, \underbrace{y_2, z_2}\ s_1$.

When $\lambda_1 = \lambda_2 = s_1$, we can find a special walk whose edges are labeled with $\underbrace{s_1, y_1}, \underbrace{z_2, y_2}, \underbrace{y_1, s_1}, \underbrace{y_2, z_2}, s_1$.

Case 2. Assume that $[w_0] = \{x_1, y_2\}$, $[w_9] = \{x_1, z_2\}$ and $\{x_1, y_2, z_2, s_1, s_2\} = \{1, 2, 3, 4, 5\}$.

When $\lambda_1 = z_2$ and $\lambda_2 = y_2$, we can find a special walk whose edges are labeled with $\underbrace{z_2, x_1}, \underbrace{s_2, z_2}, \underbrace{x_1, y_2}, \underbrace{s_1, x_1}, y_2$.

When $\lambda_1 = z_2$ and $\lambda_2 = s_2$, we can find a special walk whose edges are labeled with $\underbrace{z_2, y_2}, \underbrace{s_2, x_1}, \underbrace{y_2, s_2}, \underbrace{s_1, z_2}, s_2$.

When $\lambda_1 = s_1$ and $\lambda_2 = s_2$, we can find a special walk whose edges are labeled with $\underbrace{s_1, x_1}, \underbrace{s_2, s_1}, \underbrace{x_1, s_2}, \underbrace{s_1, x_1}, s_2$.

When $\lambda_1 = \lambda_2 = s_1$, we can find a special walk whose edges are labeled with $\underbrace{s_1, x_1}, \underbrace{z_2, y_2}, \underbrace{s_2, s_1}, \underbrace{y_2, z_2}, s_1$.

Case 3. Assume that $[w_0] = [w_9] = \{x_1, x_2\}$ and $\{x_1, x_2, s_1, s_2, s_3\} = \{1, 2, 3, 4, 5\}$.

When $\lambda_1 = s_1$ and $\lambda_2 = s_3$, we can find a special walk whose edges are labeled with $\underbrace{s_1, x_1}, \underbrace{s_3, x_2}, \underbrace{x_1, s_3}, \underbrace{s_2, x_1}, s_3$.

When $\lambda_1 = \lambda_2 = s_3$, we can find a special walk whose edges are labeled with $\underbrace{s_3, x_1}, \underbrace{s_2, x_2}, \underbrace{x_1, s_3}, \underbrace{s_1, x_1}, s_3$. This completes the proof of the theorem. □

By induction on the length of the special walk, we can conclude the following results.

Theorem 2.6. Given two vertices $w_0, w_{\ell+1}$ (not necessarily distinct) in O_n ($n \geqslant 4$) with $\lambda_1 \notin [w_0]$ and $\lambda_2 \notin [w_{\ell+1}]$, there exists an $(\ell+1)$-special walk from w_0 to $w_{\ell+1}$ such that its first edge is labeled with λ_1 and the last edge is labeled with λ_2, where ℓ is an arbitrary $(2n-1)^+$ integer.

Theorem 2.7. Given two vertices $w_0, w_{\ell+1}$ (not necessarily distinct) in O_3 with $\lambda_1 \notin [w_0]$ and $\lambda_2 \notin [w_{\ell+1}]$, there exists an $(\ell+1)$-special walk from w_0 to $w_{\ell+1}$ such that its first edge is labeled with λ_1 and the last edge is labeled with λ_2, where ℓ is an arbitrary 8^+ integer.

In the above theorems, we require the first and the last edges are prescribed, but they can also be stated in an avoiding way.

Theorem 2.8. Given two vertices w_1, w_ℓ (not necessarily distinct) in O_n ($n \geqslant 4$) with $\lambda_1 \notin [w_1]$ and $\lambda_2 \notin [w_\ell]$, there exists an $(\ell-1)$-special walk from w_1 to w_ℓ such that its first edge is not labeled with λ_1 and the last edge is not labeled with λ_2, where ℓ is an arbitrary $(2n-1)^+$ integer.

Theorem 2.9. Given two vertices w_1, w_ℓ (not necessarily distinct) in O_3 with $\lambda_1 \notin [w_1]$ and $\lambda_2 \notin [w_\ell]$, there exists an $(\ell-1)$-special walk from w_1 to w_ℓ such that its first edge is not labeled with λ_1 and the last edge is not labeled with λ_2, where ℓ is an arbitrary 8^+ integer.

2.2 Sharpness of the length in Theorem 2.6 and Theorem 2.7

Theorem 2.10. The length of the special walk in Theorem 2.6 is best possible.

Proof. Let $[w_0] = \{y_1, y_2, \ldots, y_{n-1}\}$, $[w] = \{z_1, z_2, \ldots, z_{n-1}\}$ and $\lambda_1 = s_1$. By Theorem 2.1, if we go from w_0 to w such that the first edge is labeled with λ_1, then the length of a shortest even path is at least $2n$. Hence, the length of the special walk in Theorem 2.6 cannot be improved to a smaller even number.

And if $[w_{2n}] = \{y_1, y_2, \ldots, y_{n-1}\}$, we go from w_0 to w_{2n} such that both the first edge and the last edge are labeled with s_1, in other words, we go from w_0 to w such that the first edge is labeled with s_1, and then go through the edge ww_{2n}, then the length of a shortest odd special walk from w_0 to w_{2n} is at least $2n+1$. Hence, the length of the special walk in Theorem 2.6 cannot be improved to a smaller odd number. □

Theorem 2.11. The length of the special walk in Theorem 2.7 is best possible.

Proof. Firstly, we show that the length 6 is impossible. For example, let $[w_0] = [u_0] = \{1,2\}$ and $[w_1] = [u_1] = \{3,4\}$, if we want to go from w_0 to u_0 such that the first edge is w_0u_0 and the last edge is u_1u_0, then we should go from w_1 to u_1 by a special walk of length 4, a contradiction.

Secondly, we show that the length 7 is impossible. For example, let $[w_0] = [w_7] = \{1,2\}$, $[w_1] = \{3,4\}$ and $[w_6] = \{4,5\}$. Suppose that there exists a 7-special walk $w_0w_1w_2w_3w_4w_5w_6w_7$. Note that w_1w_2 can be labeled with 1 or 2, we may assume that it is labeled with 1, thus $[w_2] = \{2,5\}$. We may assume that $w_2w_3, w_3w_4, w_4w_5, w_5w_6$ are labeled with $\lambda_3, \lambda_4, \lambda_5, \lambda_6$, respectively. If $5 \notin [w_4]$, then we should go along $w_4w_3w_2$ such that $[w_2] = \{2,5\}$, and then $\lambda_4 = 5$; similarly, we should go along $w_4w_5w_6$ such that $[w_6] = \{4,5\}$, and again we have that $\lambda_5 = 5$, a contradiction. So we have that $[w_4] = \{5, \lambda\}$. Notice that $\lambda_3 \in [w_4]$ and $\lambda_6 \in [w_4]$, thus $\lambda = \lambda_3 = \lambda_6$, but this is impossible since $\lambda_3 \in \{3,4\}$ and $\lambda_6 \in \{1,2\}$.

Thirdly, we show that the length 8 is impossible. For example, let $[w_0] = [w_7] = \{1,2\}$ and $[w_1] = [w_8] = \{3,4\}$. Suppose that there exists a 8-special walk $w_0w_1w_2w_3w_4w_5w_6w_7w_8$ and $w_1w_2, w_2w_3, w_3w_4, w_4w_5, w_5w_6, w_6w_7$ are labeled with $\lambda_2, \lambda_3, \lambda_4, \lambda_5, \lambda_6, \lambda_7$ respectively. By Theorem 2.2, the three elements $\lambda_2, \lambda_4, \lambda_6$ are distinct and $\{1,2\} \subseteq \{\lambda_2, \lambda_4, \lambda_6\}$. So we may assume that $\{\lambda_2, \lambda_4, \lambda_6\} = \{1,2,\lambda\}$. Note that both w_0w_1 and w_7w_8 are labeled with 5, thus $\lambda \neq 5$ and $\lambda \in \{3,4\}$. Note that $\lambda_2 \notin \{3,4\}$, so by Theorem 2.2, we have that $\lambda = \lambda_4 \in \{3,4\}$. By the construction of the odd graph, we have that $\lambda \notin [w_3]$, but $[w_3]$ is obtained from $\{3,4\}$ by replacing λ_3 with λ_2, and then $\lambda_3 = \lambda = \lambda_4$, a contradiction. □

3 Its applications on strong edge coloring

An ℓ-thread in a graph G is a path of length $\ell+1$ in G whose ℓ internal vertices have degree 2 in the full graph G. Note that the ends of the ℓ-thread can be the same vertex. For example, an $(\ell+1)$-cycle contains many ℓ-threads.

Let R be a subgraph of G. We call R a *κ-reducible configuration* if there exists a subgraph R' of R such that each strong κ-edge-coloring of $G - R'$ can be extended to a strong κ-edge-coloring of G.

Now, we address the following result for reducible configurations.

Theorem 3.1. Let G be a graph with maximum degree at most κ.

1. If $\kappa \geqslant 4$, then $(2\kappa-1)^+$-caterpillar is $(2\kappa-1)$-reducible. Moreover, $(2\kappa-2)$-caterpillar is not $(2\kappa-1)$-reducible.
2. If $\kappa = 3$, then 8^+-caterpillar is 5-reducible. Moreover, 7-caterpillar is not 5-reducible.

Proof. Suppose that G has a caterpillar tree with central path $u_0u_1u_2\ldots u_\ell u_{\ell+1}$, where $\ell \geqslant 2\kappa-1$ if $\kappa \geqslant 4$, and $\ell \geqslant 8$ if $\kappa = 3$. That is, every vertex u_i with $1 \leqslant i \leqslant \ell$ is adjacent to at least $\deg(u_i) - 2$ pendent vertices. Let R be the graph obtained from $G - \{u_2, u_3, \ldots, u_{\ell-1}\}$ by deleting all the 1-vertices adjacent to u_1 and u_ℓ, except u_0 and $u_{\ell+1}$. For any arbitrary strong edge coloring φ of R using colors from $\{1,2,\ldots,2\kappa-1\}$, we assume the color set for the edges incident with u_0 and $u_{\ell+1}$ is a subset of a κ-set A and a κ-set B, respectively. By Theorem 2.6 and 2.7, we can find a special walk $w_0w_1\ldots w_\ell w_{\ell+1}$ in the odd graph O_κ such that it satisfies the following properties:

(a) its length is exactly $\ell+1$;

(b) $[w_0] = \overline{A}$ and $[w_{\ell+1}] = \overline{B}$.

(c) w_0w_1 is labeled with $\phi(u_0u_1)$ and $w_\ell w_{\ell+1}$ is labeled with $\phi(u_\ell u_{\ell+1})$.

Hence, we can color the edges incident to u_i with colors in $\overline{[w_i]}$, and the edge u_iu_{i+1} is colored with the label on w_iw_{i+1} for $1 \leqslant i \leqslant \ell$. Finally, we obtain a strong edge coloring of G using colors from $\{1,2,\ldots,2\kappa-1\}$, a contradiction. □

Before we prove the main theorems, we need the following result due to Nešetřil, Raspaud and Sopena [24].

Theorem 3.2 (Nešetřil, Raspaud and Sopena [24])**.** If G is a planar graph with girth at least $5\ell+1$, then G contains a 1-vertex or an ℓ-thread.

Theorem 3.3. If G is a planar graph with maximum degree at most Δ and girth at least $10\Delta-4$, where $\Delta \geqslant 4$, then G has a strong edge coloring with $2\Delta-1$ colors.

Proof. Let G be a counterexample to the theorem with minimum number of vertices. It is obvious that G is connected and the minimum degree is at least one. Let A be the set of vertices with degree one in G, and let $H = G - A$. Note that H is connected and $\delta(H) \geqslant 1$, for otherwise G is a star and $\chi'_s(G) = \Delta(G) < 2\Delta-1$.

Claim 1. The minimum degree of H is at least two.

Proof. Suppose that xy is an edge in H with $\deg_H(x) = 1$. Thus, the vertex x is adjacent to a 1-vertex z in G. By the minimality of G, the graph $G-z$ has a strong edge coloring with $2\Delta-1$ colors. Note that xz has at most $(\deg_G(x)-2)+(\deg_G(y)-1)$ edges within distance two, so we can extend the strong edge coloring of $G-z$ to G. □

By Theorem 3.2, the graph H has a $(2\Delta-1)$-thread. Correspondingly, G has a $(2\Delta-1)$-caterpillar tree, this is contradiction due to Theorem 3.1. □

Using the same techniques, we can prove the following result which has been proved by Borodin and Ivanova [3].

Theorem 3.4 (Borodin and Ivanova [3])**.** If G is a subcubic planar graph with girth at least 41, then G has a strong edge coloring with 5 colors.

Next, we give some results with the condition on maximum average degree.

Theorem 3.5 (Cranston and West [8])**.** If the average degree of G is less than $2+\frac{2}{3\ell-1}$ and G has no 2-regular component, then G contains a 1^--vertex or an ℓ-thread.

Chang et al. [6] gave the follow result for closed special walks.

Theorem 3.6 (Chang et al. [6])**.** Every vertex is contained in a closed 2ℓ-special walk in O_Δ for arbitrary integers $\Delta, \ell \geqslant 3$.

Let $C_{\kappa,\Delta}$ be the graph obtained from C_κ by adding $(\Delta-2)$ pendant edges to each vertex in C_κ. We can easily address the following result.

Theorem 3.7. If κ is even and at least 6, or κ is odd and at least $2\Delta-1$ with $\Delta \geqslant 4$, then $\chi'_s(C_{\kappa,\Delta}) = 2\Delta-1$.

Theorem 3.8. Let G be a graph with maximum degree at most Δ ($\Delta \geqslant 4$). If $\text{evengirth}(G) \geqslant 6$, $\text{oddgirth}(G) \geqslant 2\Delta-1$ and $\text{mad}(G) < 2+\frac{1}{3\Delta-2}$, then $\chi'_s(G) \leqslant 2\Delta-1$.

Proof. Let G be a counterexample to the theorem with minimum number of vertices. It is obvious that G is connected and the minimum degree is at least one. Let A be the set of vertices with degree one in G, and let $H = G-A$. It is easy to show that the minimum degree of H is at least two.

If H has a $(2\Delta-1)$-thread, then G has a $(2\Delta-1)$-caterpillar tree, which contradicts Theorem 3.1. So we may assume that there exists no $(2\Delta-1)$-threads in H. Note that H is connected. By Theorem 3.5 and the hypothesis, G is a subgraph of $C_{\kappa,\Delta}$, this is a contradiction due to Theorem 3.7. □

Similarly, we have the following result for subcubic graphs due to Theorem 2.5, 3.5 and 3.7.

Theorem 3.9. If G is a subcubic graph with girth at least 8 and $\text{mad}(G) < 2+\frac{2}{23}$, then $\chi'_s(G) \leqslant 5$.

Remark 1. The essential of all the proofs is finding a special walk with fixed length between any two vertices in the odd graph, in which the first edge and the last edge are specified. Obviously, all the results cannot be directly extended to the list version of strong edge coloring.

Acknowledgments. This project was supported by the National Natural Science Foundation of China (11101125) and partially supported by the Fundamental Research Funds for Universities in Henan.

References

[1] L. D. Andersen, The strong chromatic index of a cubic graph is at most 10, Discrete Math. 108 (1-3) (1992) 231–252.

[2] C. L. Barrett, G. Istrate, V. S. Anil Kumar, M. V. Marathe, S. Thite and S. Thulasidasan, Strong edge coloring for channel assignment in wireless radio networks, in: Proceedings of the Fourth Annual IEEE International Conference on Pervasive Computing and Communications Workshops, 2006, pp. 106–110.

[3] O. V. Borodin and A. O. Ivanova, Precise upper bound for the strong edge chromatic number of sparse planar graphs, Discuss. Math. Graph Theory 33 (4) (2013) 759–770.

[4] H. Bruhn and F. Joos, A stronger bound for the strong chromatic index, (2015) arXiv:1504.02583.

[5] G. J. Chang and D. D.-F. Liu, Strong edge-coloring for cubic Halin graphs, Discrete Math. 312 (8) (2012) 1468–1475.

[6] G. J. Chang, M. Montassier, A. Pêcher and A. Raspaud, Strong chromatic index of planar graphs with large girth, Discuss. Math. Graph Theory 34 (4) (2014) 723–733.

[7] G. J. Chang and N. Narayanan, Strong chromatic index of 2-degenerate graphs, J. Graph Theory 73 (2) (2013) 119–126.

[8] D. W. Cranston and D. B. West, A guide to the discharging method, (2013) arXiv:1306.4434.

[9] M. Dębski, J. Grytczuk and M. Śleszyńska-Nowak, The strong chromatic index of sparse graphs, Inform. Process. Lett. 115 (2) (2015) 326–330.

[10] P. Erdős, Problems and results in combinatorial analysis and graph theory, Discrete Math. 72 (1-3) (1988) 81–92.

[11] R. J. Faudree, A. Gyárfás, R. H. Schelp and Z. Tuza, Induced matchings in bipartite graphs, Discrete Math. 78 (1-2) (1989) 83–87.

[12] R. J. Faudree, R. H. Schelp, A. Gyárfás and Z. Tuza, The strong chromatic index of graphs, Ars Combin. 29 (B) (1990) 205–211.

[13] J.-L. Fouquet and J.-L. Jolivet, Strong edge-colorings of graphs and applications to multi-k-gons, Ars Combin. 16 (A) (1983) 141–150.

[14] J.-L. Fouquet and J.-L. Jolivet, Strong edge-coloring of cubic planar graphs, in: Progress in graph theory (Waterloo, Ont., 1982), Academic Press, Toronto, ON, 1984, pp. 247–264.

[15] Z. Guo, X. Zhao and Y. Yi, Strong edge coloring of planar graphs using $2\Delta - 1$ colors, (2015+) submitted.

[16] P. Horák, H. Qing and W. T. Trotter, Induced matchings in cubic graphs, J. Graph Theory 17 (2) (1993) 151–160.

[17] A. V. Kostochka, X. Li, W. Ruksasakchai, M. Santana, T. Wang and G. Yu, Strong chromatic index of subcubic planar multigraphs, European J. Combin. 51 (2015) 380–397.

[18] H.-H. Lai, K.-W. Lih and P.-Y. Tsai, The strong chromatic index of Halin graphs, Discrete Math. 312 (9) (2012) 1536–1541.

[19] K.-W. Lih and D. D.-F. Liu, On the strong chromatic index of cubic Halin graphs, Appl. Math. Lett. 25 (5) (2012) 898–901.

[20] M. Molloy and B. Reed, A bound on the strong chromatic index of a graph, J. Combin. Theory Ser. B 69 (2) (1997) 103–109.

[21] K. Nakprasit, A note on the strong chromatic index of bipartite graphs, Discrete Math. 308 (16) (2008) 3726–3728.

[22] K. Nakprasit and K. Nakprasit, The strong chromatic index of graphs and subdivisions, Discrete Math. 317 (2014) 75–78.

[23] T. Nandagopal, T.-e. Kim, X. Gao and V. Bharghavan, Achieving MAC layer fairness in wireless packet networks, in: MobiCom '00: Proceedings of the 6th annual international conference on Mobile computing and networking, 2000, pp. 87–98.

[24] J. Nešetřil, A. Raspaud and E. Sopena, Colorings and girth of oriented planar graphs, Discrete Math. 165/166 (1997) 519–530.

[25] S. Ramanathan, A unified framework and algorithm for channel assignment in wireless networks, Wireless Networks 5 (2) (1999) 81–94.

[26] W. C. Shiu, P. C. B. Lam and W. K. Tam, On strong chromatic index of Halin graph, J. Combin. Math. Combin. Comput. 57 (2006) 211–222.

[27] W. C. Shiu and W. K. Tam, The strong chromatic index of complete cubic Halin graphs, Appl. Math. Lett. 22 (5) (2009) 754–758.

[28] A. Steger and M.-L. Yu, On induced matchings, Discrete Math. 120 (1-3) (1993) 291–295.

[29] T. Wang, Strong chromatic index of k-degenerate graphs, Discrete Math. 330 (2014) 17–19.

[30] J. Wu and W. Lin, The strong chromatic index of a class of graphs, Discrete Math. 308 (24) (2008) 6254–6261.

[31] G. Yu, Strong edge-colorings for k-degenerate graphs, Graphs Combin. (2014) doi:10.1007/s00373–014–1478–1.